

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : G01N 33/543, 33/00, C12N 5/00 C12N 1/20</p>	<p>A1</p>	<p>(11) International Publication Number: WO 90/10228 (43) International Publication Date: 7 September 1990 (07.09.90)</p>
<p>(21) International Application Number: PCT/US90/01060 (22) International Filing Date: 1 March 1990 (01.03.90) (30) Priority data: 317,407 1 March 1989 (01.03.89) US Not furnished 26 February 1990 (26.02.90) US (71) Applicant: THE UNITED STATES OF AMERICA, represented by THE SECRETARY, UNITED STATES DEPARTMENT OF COMMERCE [US/US]; Washington, DC 20231 (US). (72) Inventors: LIOTTA, Lance, A. ; 9027 Mistwood Drive, Potomac, MD 20854 (US). STETLER-STEVENSON, William ; 11227 White Barn Court, Gaithersburg, MD 20879 (US). KRUTZSH, Henry ; 9704 DePaul Drive, Bethesda, MD 20817 (US).</p>		<p>(74) Agents: OLIFF, James, A. et al.; Oliff & Berridge, 277 S. Washington Street, Alexandria, VA 22314 (US). (81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published With international search report.</p>

(54) Title: MATRIX METALLOPROTEINASE PEPTIDES: ROLE IN DIAGNOSIS AND THERAPY

(57) Abstract

A family of metalloproteinases exist which cleave extracellular matrix molecules. These metalloproteinases are secreted in a latent inactive form and require activation in order to specifically cleave the preferred substrate. A series of peptides have been prepared based on the complete sequence analysis of type IV procollagenase. Peptide inhibitors were synthesized which correspond to cysteine repeat regions and histidine containing regions; the mechanism of action of these peptides involves inhibition of binding of the enzyme to the substrate. Peptide inhibitors were synthesized which correspond to the peptide cleaved off during activation, and constitute a novel class of metalloproteinase inhibitors. These inhibitors are members of a series of peptides which contain the core amino acid sequence RKPRC or analogs thereof. The cysteine residue is required for activity. Affinity purified antibodies directed against specific peptides can be used to a) detect any general metalloproteinase enzyme with the sequence in part VAAHE or PRCGNPD, and distinguish it from other known members of the metalloproteinase family, b) block functional domains resulting in the inhibition of enzyme activity, and c) distinguish latent from activated forms of the enzyme.

inhibitor MW

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Malì
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

- 1 -

MATRIX METALLOPROTEINASE PEPTIDES:

ROLE IN DIAGNOSIS AND THERAPY

BACKGROUND OF THE INVENTIONField of the invention

5 This invention relates to peptides useful in metalloproteinase detection and inhibition. Specifically, the invention relates to peptides derived from the sequence of type IV collagenase which correspond to domains of the enzyme which are involved in activation of the enzyme and interaction of the enzyme with its sub-strate. Antibodies recognizing the peptides are useful in enzyme detection. Specific peptides, identified by means of functional studies, constitute new classes of metalloproteinase inhibitors.

Background

15 The degradation of interstitial and basement membrane collagens is initiated by a specific class of metalloproteinase, the matrix metalloproteinases, also known as the collagenases (EC 3.4.24.7), which are secreted into the extracellular matrix in zymogen form. Members of this collagenase gene family include: the interstitial collagenases, which degrade collagen types I, II and III and have been characterized with respect to substrate specificity and requirements for activation (Stricklin, G.P., Jeffrey, J.J., Rosewit, W.T., and Eisen, A.Z., 1983, Biochemistry 22, 61-68; Goldberg, G.I., Wilhlem, S., Kronberger, A., Bauer, E.A., Grant, G.A., and Eisen, A.Z., 1986, J. Biol. Chem. 261, 6600-6605; Hasty, K.A., Jeffrey, J.J., Hibbs, M.S., and Welgus, H.G., 1987, J. Biol. Chem. 262, 10048-1052; Fields, G.B., Van Wart, H.E., and Birkedal-Hansen, H., 1987, J. Biol. Chem. 262, 6221-6226; Grant, G.A., Eisen, A.Z., Marmer, B.L. Rosweit, W.T., and Goldberg, G.I., 1987, J. Biol. Chem. 262, 5886-5889); stromelysin, which degrades proteoglycans, glycoproteins, and the non-helical portions of collagenous molecules (Wilhelm,

- 2 -

S.M., Collier, I.E., Kronberger, A., Eisen, A.Z., Marmer, B.L., Grant, G.A., Bauer, E., and Goldberg, G.I., 1987, Proc. Natl. Acad. Sci. U.S.A. **84**, 6725-6729; Whitman, S.E., Murphy, G., Angel, P., Rahmsdorf, H.-J., Smith, B.J., Lyons, A., Harris, T.J.T., Reynolds, J.J., Herrlich, P. and Docherty, A.J.P., 1986, Biochem. J. **240**, 913-916); and type IV collagenase, which degrades pepsin-resistant triple-helical type IV collagen and interstitial collagens (gelatin). Type IV collagenase has been identified in human tumor cells (Liotta, L.A., Kleinerman, J., Catanzaro, P., and Rynbrandt, D., 1977, J. Natl. Cancer Inst. **58**, 1427-1439; Turpeenniemi-Hujanen, T., and Tryggvason, K., 1982, Int. J. Cancer **30p**, 669-673; Liotta, L.A., Abe, S., Gehron-Robey, P., and Martin, G.R., 1979, Proc. Natl. Acad. Sci. U.S.A. **76** 2268-2272; Liotta, L.A., Tryggvasson, K., Garbisa, S., Hart, I., Foltz, C.M., and Shafie, S., 1980, Nature, Lond., **284**, 67-68; Collier, I.E., Wilhelm, S.M., Eisen, A.Z., Marmer, B.L., Grant, G.A., Seltzer, J.L., Kronberger, A., He, C., Bauer, E.A., and Goldberg, G.I., 1988, J. Biol. Chem. **263**, 6579-6587), endothelial cells (Kalebic, T., Barbisa, S., Glaser, B., and Liotta, L.A., 1983, Science **221**, 281-283), bone (Murphy, G., McAlpine, C.G., Poll, C.T., and Reynolds, J.J., 1985, Biochem. Biophys. Acta **831**, 49-58), fibroblasts (Collier, I.E., Wilhelm, S.M., Eisen, A.Z., Marmer, B.L., Grant, G.A., Seltzer, J.L., Kronberger, A., He, C., Bauer, E.A., and Goldberg, G.I., 1988, J. Biol. Chem. **263**, 6579-6587), polymorphonuclear leukocytes (Uitto, V.J., Schwartz, D., and Veis, A., 1980, Eur. J. Biochem. **105**, 409-417) and macrophages (Garbisa, S., Ballin, M., Daga-Giordini, D., Fastelli, G., Natural, M., Negro, A., Semenzato, G., and Liotta, L.A., 1986, J. Biol. Chem. **261**, 2369-2375). This enzyme is a neutral metalloproteinase of 68 to 72 kilodaltons which is secreted in

- 3 -

zymogen form (Liotta, L.A., Abe, S., Gehron-Robey, P., and Martin, G.R., 1979, Proc. Natl. Acad. Sci. U.S.A. 76, 2268-2272; Liotta, L.A., Tryggvasson, K., Garbisa, S., Gehron-Robey, P., and Abe, S., 1981, Biochemistry 20, 100-104; Salo, T., Liotta, L.A., and Tryggvasson, K., 1983, J. Biol. Chem. 258, 3058-3063). In addition, several other members of this collagenase gene family have been described recently, including a second type of stromelysin (stromelysin-2), a 92 kilodalton form of type IV collagenase, and Putative Uterine Metallo-proteinase (PUMP)-1, a low molecular weight uterine collagenase (Wilhelm, S.M., Collier, I.E., Marmer, B.L., Eisen, A.Z., Grant, G.A., and Goldberg, G.I., 1989, J. Biol. Chem. 264, 17213-17221; Woessner, J. F. and Talpin, C. J., 1988, J. Biol. Chem. 263, 16918-16925).

The 70 kilodalton type IV collagenase has been closely linked to the metastatic potential of tumors in murine tumor models (Liotta, L.A., Tryggvasson, K., Garbisa, S., Hart, I., Foltz, C.M., and Shafie, S., 1980, Nature, Lond, 284, 67-68) and is augmented following H-ras oncogene induced genetic induction of the metastatic phenotype (Muschel, R., Williams, J.E., Lowy, D.R., and Liotta, L.A., 1985, Amer. J. Pathol. 121, 1-8; Garbisa, S., Pozzatti, R., Muschel, R.J., Saffiotti, U., Ballin, M., Goldfarb, R.H., Khoury, G., and Liotta, L.A., 1987, Cancer Res. 47, 1523-1528). Trypsin treatment results in activation of the latent enzyme and a concomitant reduction in the molecular mass (Liotta, L.A., Tryggvasson, K., Garbisa, S., Gehron-Robey, P., and Abe, S., 1981, Biochemistry 20, 100-104). Organomercurial compounds have also been shown to activate this enzyme, and these are also associated with a reduction in the molecular mass (Collier, I.E., Wilhelm, S.M., Eisen, A.Z., Marmer, B.L., Grant, G.A., Seltzer, J.L., Kronberger, A., He, C., Bauer, E.A., and Goldberg,

- 4 -

5 G.I., 1988, J. Biol. Chem. 263, 6579-6587; Murphy, G.,
McAlpine, C.G., Poll, C.T., and Reynolds, J.J., 1985,
Biochem. Biophys. Acta 831, 49-58). The activated
enzyme cleaves type IV collagen to generate charac-
10 teristic 1/4 amino-terminal and 3/4 carboxy-terminal
fragments (Liotta, L.A., Tryggvasson, K., Garbis, S.,
Gehron-Robey, P., and Abe, S., 1981, Biochemistry 20,
100-104; Collier, I.E., Wilhelm, S.M., Eisen, A.Z.,
Marmer, B.L., Grant, G.A., Seltzer, J.L., Kronberger,
15 A., He., C., Bauer, E.A., and Goldberg, G.I., 1988, J.
Biol. Chem. 263, 6579-6587; Fessler, L.I., Duncan,
K.G., Fessler, J.H., Salo, T., and Tryggvason, K.,
1984, J. Biol. Chem. 259, 9783-9789). It has also
been demonstrated that gelatinolytic activity is
15 associated with this enzyme (Collier, I.E., Wilhelm,
S.M., Eisen, A.Z., Marmer, B.L., Grant, G.A., Seltzer,
J.L., Kronberger, A., He., C., Bauer, E.A., and
Goldberg, G.I., 1988, J. Biol. Chem. 263, 6579-6587;
Hoyhtya, M., Turpeenniemi-Hujanen, T., Stetler-
20 Stevenson, W., Kruttsch, H., Tryggvason, K., and
Liotta, L.A., 1988, FEBS Letters 233, 109-113; Murphy,
G., McAlpine, C.G., Poll, C.T., and Reynolds, J.J.,
1985, Biochem. Biophys. Acta 831, 49-58) as well as a
type V collagenolytic activity (Collier, I.E.,
25 Wilhelm, S.M., Eisen, A.Z., Marmer, B.L., Grant, G.A.,
Seltzer, J.L., Kronberger, A., He., C., Bauer, E.A.,
and Goldberg, G.I., 1988, J. Biol. Chem. 263, 6579-
6587; Murphy, G., McAlpine, C.G., Poll, C.T., and
Reynolds, J.J., 1985, Biochem. Biophys. Acta 831, 49-
30 58).

Type IV collagenase has been purified from
human melanoma cells and sequence information on the
intact protein amino terminus has been obtained as
well as on tryptic and cyanogen bromide peptide
35 fragments (Hoyhtya, M., Turpeenniemi-Hujanen, T.,
Stetler-Stevenson, W., Kruttsch, H., Tryggvason, K.,
and Liotta, L.A., 1988, FEBS Letters 233, 109-113).

- 5 -

Th sequence informati n demonstrates that type IV collagenas shows limit d s qu nc homology to interstitial collagenas and stromelysin. A recent report has characterized a partial cDNA clone for a metalloproteinase secreted by H-ras-transformed human bronchial epithelial cells (Collier, I.E., Wilhelm, S.M., Eisen, A.Z., Marmer, B.L., Grant, G.A., Seltzer, J.L., Kronberger, A., He., C., Bauer, E.A., and Goldberg, G.I., 1988, J. Biol. Chem. **263**, 6579-6587). The transformed bronchial epithelial enzyme is capable of specifically degrading type IV collagen, and the deduced amino acid sequence shows identity with that reported for tryptic and cyanogen bromide fragments of human tumor IV collagenase (Hoyhtya, M., Turpeenniemi-Hujanen, T., Stetler-Stevenson, W., Kruttsch, H., Tryggvason, K., and Liotta, L.A., 1988, FEBS Letters **233**, 109-113). Thus, human melanoma cell type IV collagenase appears identical with the enzyme from H-ras-transformed bronchial epithelial cells, which is also found in fibroblasts (Collier, I.E., Wilhelm, S.M., Eisen, A.Z., Marmer, B.L., Grant, G.A., Seltzer, J.L., Kronberger, A., He., C., Bauer, E.A., and Goldberg, G.I., 1988, J. Biol. Chem. **263**, 6579-6587) and bone cell explants (Murphy, G., McAlpine, C.G., Poll, C.T., and Reynolds, J.J., 1985, Biochem. Biophys. Acta **831**, 49-58).

SUMMARY OF THE INVENTION

The complete amino acid sequence of the zymogen form of the type IV collagenase purified from human melanoma cells has now been analyzed and a series of inhibitory synthetic peptides corresponding to a variety of domains of the enzyme have been prepared. These domains include: the 80 residue amino terminus which is cleaved from the proenzyme during activation; a cystein -rich interior domain; a histidine-containing r gion; and a region 159 residues from th carboxy terminus. P ptides from these

- 6 -

domains have been used to generate antibodies against specific domains within the type IV collagenase molecule. In the present invention, the antibodies, direct amino acid sequence analysis, and peptides have been used to determine a) the region of the enzyme involved in binding and interaction with the substrate, and b) the structure of the major type IV collagenase conversion product produced during zymogen activation with the organomercurial compound p-amino-phenylmercuric acetate.

It was discovered that type IV collagenase proenzyme activation by an organomercurial compound, PAPMA, is accompanied by an autocatalytic removal of an 80 amino acid amino terminal fragment, resulting in a stable, active enzyme species of 62 kDa. Furthermore, these data show that type IV collagenase shares not only sequence homology but functional domain identity with the amino terminal regions of other extracellular matrix degrading metalloproteinases.

As shown in Figures 8 and 9, below, an amino terminal peptide (residues 1-80) is cleaved off during type IV collagenase activation. This discovery raised the possibility that this peptide contains an intrinsic enzyme inhibitor which blocks the active site and renders the enzyme inactive. Removal of this amino terminal segment during activation thus would remove this inhibitor and exposes the active site. The critical region involved in the inhibition is the region enclosed in a box in Figure 10 which contains an unpaired cysteine residue. This region shows a conserved nature in the other matrix metalloproteinases, and it has a high probability of beta turn conformation by Chou Fassman analysis. Furthermore, it was reasonable to hypothesize that the unpaired cysteine residue in this sequence interacted in a non-covalent fashion with the metal ion in the active site of the enzyme. Organomercurial activation (it is

- 7 -

known that APMA binds to sulfhydryl residues) would thus disrupt this interaction and cause a conformational change which would separate the inhibitor segment from proximity to the active site. To test his completely novel hypothesis, synthetic peptides were prepared which correspond to a series of overlapping regions in the amino terminal residues 1-87. As shown in Figures 11 and 12 and Table 4, below, only those peptides incorporating the conserved region containing the unpaired cysteine were strongly inhibitory at concentration less than 0.1 mM. The cysteine was required for the activity. These peptides, therefore, constitute a highly novel class of inhibitors for metalloproteinases.

In summary, the discoveries stemming from this analysis have led to identification of a) a region near the amino terminus of type IV procollagenase that constitutes an intrinsic enzyme inhibitor which may block the active site of the enzyme when the enzyme is in a latent state, and b) a region near the middle of the enzyme that is involved in binding and interaction with the substrate. Peptides homologous to these regions constitute novel matrix metalloproteinase inhibitors.

DESCRIPTION OF THE DRAWINGS

FIGURE 1 illustrates the complete amino acid sequence of human type IV procollagenase secreted by tumor cells.

FIGURE 2 illustrates a comparison of amino acid homologies between type IV procollagenase and prostromelysin and procollagenase I organized into domains.

FIGURE 3 (A) ELISA characterization of anti-peptide antibody A1-17. The synthetic peptide corresponding to the amino terminal residues 1 through 17 of type IV procollagenase was synthesized and used

- 8 -

an antigen. The peptide-bovine serum albumin conjugate was used as coating antigen in this ELISA.

(B) ELISA characterization of antipeptide antibody A472-490. The synthetic peptide corresponding to the internal residues 472-490 of type IV procollagenase was synthesized and used as an antigen. The peptide-bovine serum albumin conjugate was used as coating antigen in this ELISA.

(C) Competition ELISA assays were performed using the appropriate peptide for each antibody. Peptide-bovine serum albumin was used as coating antigen and free peptides were used as competing antigens.

(D) Western blots of crude and gelatin-affinity purified type IV procollagenase. Shown are crude type IV procollagenase immunoblotted with A1-17 (20 μ l of A2058 melanoma cell conditioned media, lane a), purified type IV collagenase immunoblotted with A472-490 (30 ng, lane c).

FIGURE 4 illustrates the assay for type IV collagenase binding to its substrate type IV collagen. Purified type IV collagenase was used to compete for the binding of labeled type IV collagenase to pep-sinized type IV collagen coated in microtiter wells (mean \pm S.D.). Saturation of binding is demonstrated in the lower curve using increasing amounts of substrate.

FIGURE 5 illustrates the gelatin zymogram inhibition of type IV collagenase by the designated synthetic peptides. The gelatinase activity is visualized as white cleared bands at approximately 70 kDa. The white cleared bands are abolished in the presence of the inhibitor. Three replicates are shown.

FIGURE 6 illustrates an example inhibition of binding of labeled type IV collagenase to type IV collagen by the designated peptides. The histidine

- 9 -

containing peptid is derived from a region in fibronectin homologous to the histidine containing domain of type IV collagenase shown in Figures 2 and 5. The histidine residue is required for binding competition.

5 FIGURE 7 illustrates a time course for PAPMA activation of type IV procollagenase followed by gelatin zymogram and western blotting. 20 μ l aliquots of A2058 melanoma cell conditioned media were activated in the presence of 1.0 mM PAPMA for the indicated times (min). The reactions were stopped by the addition of EDTA to 10 mM and the samples were electrophoresed on a 9% acryl-amide gel with or without gelatin. The gelatin containing gels were developed as zymograms after electrophoresis. The nongelatin containing gels were electrophoretically transferred to nitrocellulose and then immunostained with the designated affinity purified antibodies (1 μ g/ml). Loss of the amino terminal antigenic domain occurs during the PAPMA induced conversion from the 70 kDa to the 62 kDa form. MWM, prestained molecular weight markers.

 FIGURE 8 illustrates a time course of PAPMA activation of purified type IV collagenase followed by type IV collagen degradation assay. 10 μ l aliquots of purified type IV collagenase (23 μ g/ml) were made 1 mM in PAPMA and preincubated at 37°C for the indicated times. The samples were then diluted to 60 μ l final volume by the addition of 50 mM Tris HCl, 0.15 M NaCl, 5 mM CaCl_2 , pH 7.6 ^3H -type IV collagen (New England Nuclear) was added and the reaction mix allowed to incubate at 37°C for 30 min. Samples were assayed in triplicate. Maximal activity corresponds to 2.8 μ g type IV collagen degraded/h/ μ g purified enzyme at 37°C.

35 FIGURE 9 illustrates the amino acid sequences of the latent and active forms of type IV procollagenase as determined by direct amino acid

- 10 -

s qu ncing of th purified enzyme. Also shown is th cleavage site of the autocatalysis on pAPMA activation. The insert shows the apparent molecular weights on a silver stained NaDodSO₄-PAGE gel of the purified latent (20 ng, lane A) and active enzymes (20 ng, lane B).

FIGURE 10 illustrates the amino termini (residues 1-110) of type IV procollagenase (top line), interstitial procollagenase (middle line), and prostromelysin (bottom line). The area of homology just upstream from the cleavage site is shown in box A, the cleavage sites following pAPMA activation are shown in box B and the cysteine residues are underlined. Additional reported sites of cleavage following pAPMA treatment of interstitial collagenase and stromelysin are denoted by asterisks.

FIGURE 11 illustrates the dose dependent inhibition of purified activated type IV collagenase cleavage of pepsinized type IV collagen by the designated synthetic peptides. Peptide TMRKPRCGNPDVAN at a concentration of 0.1 mM inhibits 80% of the enzyme activity. Higher concentrations abolish all enzyme activity.

FIGURE 12 illustrates a comparison of peptides tested for enzyme inhibitory activity. The peptides were derived from the amino terminal sequence shown in Figure 10. The core sequence PRCG is necessary for inhibitory activity based on the fact that peptides lacking this sequence are devoid of significant inhibitory activity. Further refinement of the minimal sequence requirements was accomplished using the peptides listed in Table 2.

FIGURE 13 illustrates the nucleotide sequenc of th portion of a human cDNA clone for human type IV procollagenas which encodes the 80 amino acid sequenc of th amino t rminal peptid that

- 11 -

is cleaved from human type IV procollagenase upon activation.

DETAILED DESCRIPTION OF THE INVENTION

5 It is the object of this invention to provide peptide sequences that have the ability to inhibit a matrix metalloproteinase.

It is a further object of the invention to provide antibodies for use in identifying the presence of a matrix metalloproteinase.

10 It is a further object of this invention to provide a method of treating patients suffering tissue damage arising from tissue destruction caused by an activated matrix metalloproteinase.

15 The complete amino acid sequence of the human type IV collagenase is illustrated in Figure 1. This sequence reveals that this protein can be divided into a series of domains as shown in Figure 2. A cysteine-rich (12 cysteine residues) domain encoded by a segment of approximately 1 kb has no significant
20 homology with other sequenced metalloproteinases such as type I collagenase and stromelysin. The cysteine-rich region, however, does show significant homology to fibronectin. The cysteine-rich sequence has never previously been directly tested for its functional
25 role in the metalloproteinase activity. Peptides derived from the cysteine-rich region (residues 200-370) inhibited enzyme substrate binding and are considered embodiments of the present invention.

30 Three other domains of type IV collagenase show significant homologies with other matrix metalloproteinases (Figure 2). In particular, a region at residues 371 to 386, designated "MBD" and illustrated in Figure 2, has closely homologous
35 sequences in all three matrix metalloproteinases including thermolysin. In the actual crystallized thermolysin, this region is related to the putative Zn binding domain of the enzyme. However, the MBD

- 12 -

sequence has never previously been directly tested for its functional role in the metalloproteinase activity. In the present invention (a) synthetic peptides derived from these regions, and (b) affinity purified antibodies directed against these regions constitute inhibitors which block gelatinase and type IV collagenase activity. Furthermore, peptides derived from fibronectin homologous to the type IV collagenase domains outlined also inhibit binding of type IV collagenase to its substrates.

The matrix metalloproteinase inhibitors of this invention can be used in the treatment of inappropriate angiogenesis, arthritis, tumor growth, invasion and metastasis, and granulomatous inflammatory conditions such as sarcoidosis. In these conditions, it is possible to estimate the amount of enzyme produced and the amount of peptide inhibitor required to inhibit greater than 90% of the active enzyme as shown in Figure 11. Therapeutic dose of the inhibitory peptide falls within an acceptable pharmacologic range of 10-250 mg/kg/d, with a more preferred dosage being 25-100 mg/kg/d. The dosage for a given patient will depend on the amount of enzyme produced in the patient, the condition and size of the patient. The inhibitors may be given as infusions or by any means which provides ready transmission into the circulation. Lyophilized powders may be "snorted". Preparations for buccal or sublingual administration may also be given. For respiratory tract involvement, the peptides may be administered by inhalation. Aerosols are particularly useful for this purpose. For conditions of the eye, the peptides may be administered as eye drops.

EXAMPLE 1

Histidine Containing Peptide Inhibitors

The synthetic peptide corresponding to residues 371-386 (the MBD region) of type IV

- 13 -

collagenase abolish d the g latinas and th collagenase type IV activity. This is demonstrated in Figure 5 using gelatin zymograms. Furthermore, affinity purified antibodies recognizing these domains also inhibited the gelatinase and collagenase type IV activities. The mechanism of action of these peptides was, at least in part, due to their ability to complete for binding of type IV collagenase to type IV collagen (Figure 6).

The metalloproteinase peptide inhibitor has a protein peptide having substantial homology with a histidine-containing domain at residues 371 to 386 of type IV collagenase. The protein peptide inhibits gelatinolytic and collagenolytic activities of metalloproteinases. The protein peptide of the metalloproteinase inhibitor preferably contains at least one histidine residue for activity. One desirable embodiment of the protein peptide of the metalloproteinase inhibitor has a histidine-containing sequence, said sequence being a member selected from the group consisting of VAAHEFGHAMGLEHSQ, VAAHEFGAAMGLEHSQ, VAAHELGHSLGLSHST, VAAHELGHSLGLSHST, VAAHEIGHSLGLFHSA, VVAHELTHAVTDYTAG and the fibronectin peptide AAHEEICTTNEGVM. Since the latter peptide was effective and required the histidine residue, the core sequence AHE was determined to be a minimum determinant of one major embodiment of the invention.

By substituting amino acids into conserved residue regions, it was demonstrated that the histidine residues were, in part, required for the inhibitory activity of the peptide as shown in Figures 5 and 6. When all three histidines were replaced with alanines, the peptide was completely inactive. However, if only the central histidine was replaced by alanine, the inhibitory activity was retained. Furthermore, replacement of the two glutamic acid (E) residues with glutamine (Q) substantially altered the

- 14 -

inhibitory activity of the peptide. These inhibitory peptides bear no homology to the cleavage site on the type IV collagen or gelatin substrate. Their mechanism of action involves an interference of the enzyme-substrate interaction in a region of the proteinase which can interact with a metal ion, such interaction being necessary for substrate cleavage. The active peptide partially inhibited the gelatinolytic activity of thermolysin under concentrations of the peptide which abolished the type IV collagenase activity. The active peptides failed to inhibit a variety of serine and thiol proteases or "non-metalloproteinases" tested, including plasmin and trypsin.

TABLE 1 is a list of the synthetic histidine-containing peptides, including amino acid substitutions, that have been tested as candidates for inhibitors of a matrix metalloproteinase, chosen from the sequence in Figure 1, based on functional studies by the inventors and based on homology comparisons with other metalloproteinases.

TABLE 1

Synthetic Histidine-Containing Peptides Tested
for Inhibition of Type-IV Collagenase (N→C)

5	1.	Val-Ala-Ala-His-Glu-Phe-Gly-His-Ala-Met-Gly-Leu-Glu-His-Ser-Gln
	2.	Val-Ala-Ala-Ala-Glu-Phe-Gly-Ala-Ala-Met-Gly-Leu-Glu-Ala-Ser-Gln
10	3.	Val-Ala-Ala-His-Gln-Phe-Gly-His-Ala-Met-Gly-Leu-Gln-His-Ser-Gln
	4.	Val-Ala-Ala-His-Glu-Phe-Gly-Ala-Ala-Met-Gly-Leu-Glu-His-Ser-Gln
15	5.	Val-Ala-Ala-Ala-Glu-Phe-Gly-His-Ala-Met-Gly-Leu-Glu-His-Ser-Gln
	6.	Val-Ala-Ala-Ala-Glu-Phe-Gly-Ala-Ala-Met-Gly-Leu-Glu-His-Ser-Gln
	7.	Val-Ala-Ala-His-Glu-Phe-Gly-Ala-Met-Gly-Leu-Glu-Ala-Ser-Gln
20	8.	Val-Ala-Ala-His-Ala-Phe-Gly-His-Ala-Gly-Leu-Ala-His-Ser-Gln
	9.	Val-Val-Ala-His-Glu-Leu-Thr-His-Ala-Val-Thr-Asp-Tyr-Thr-Ala-Gly
25	10.	Val-Ala-Ala-Glu-Lys-Phe-Gly-Glu-Ala-Met-Gly-Leu-Lys-Glu-Ser-Gln
	11.	Ala-Ala-His-Glu
	12.	Ala-Ala-His-Glu-Glu-Ile-Cys-Thr-Thr-Asn-Glu-Gly-Val-Met
30	13.	Ala-Ala-Ala-Glu-Glu-Ile-Cys-Thr-Thr-Asn-Glu-Gly-Val-Met
	14.	Thr-Met-Arg-Lys-Pro-Arg-Cys-Gly-Asn-Pro-Asp-Val-Ala-Asn-Tyr-Asn-Phe-Phe-Pro-Arg-Lys-Pro-Lys

- 16 -

TABLE 1 (continuation)

	15.	Thr-Met-Arg-Lys-Pro-Arg-Cys-Gly-Asn-Pro-Asp-Val-Ala-Asn
5	16.	Thr-Met-Arg-Lys-Pro-Arg-Ser-Gly-Asn-Pro-Asp-Val-Ala-Asn
	17.	Arg-Lys-Pro-Arg-Cys-Gly-Asn
	18.	Glu-Ser-Cys-Asn-Leu-Phe-Val-Leu-Lys-Asp-Thr-Leu-Lys-Met-Gln-Lys
10	19.	Met-Trp-Cys-Ala-Thr-Thr-Ala-Asn-Tyr-Asp-Asp-Arg-Lys-Trp-Gly-Phe-Cys-Pro-Asp-Gln-Gly-Tyr-Ser-Leu

MethodsGelatin Zymogram

15 A gelatin zymogram for visualization of gelatinase activity was prepared using the stock solutions and mixing procedures below.

Stock Solutions:Mixing Procedures:

20	a) 2M Tris HCl, pH 8.8	242 g Tris base in 800 ml of dH ₂ O, adjust pH to 8.8 with concentrated HCl and dilute to 1 liter
25	b) 0.5 M Tris pH 6.8	6.05 g Tris base in 80 ml of dH ₂ O, adjust pH to 6.8 with concentrated HCl and dilute to 100 ml
30	c) 30% Acrylamide with 0.8% bisacrylamide,	To 100 g of acrylamide, add 2.4 g of bisacrylamide, add enough dH ₂ O to dissolve acrylamide. Dilute to 333 ml and store at 4°C in an aluminum foil wrapped bottle
35	d) 10% SDS	Dissolve 100 g of SDS in dH ₂ O and dilute to 1 liter final volume
	e) 10% Ammonium persulfate	To 1 g of ammonium persulfate, add dH ₂ O to 10 ml final volume. Store at 4°C
40	f) 10 X Electrode buffer	Dissolve 144 g glycine in 600 ml of dH ₂ O, add 125 ml of 2 M Tris HCl, pH 8.8, and 100

- 17 -

- ml of 10% SDS. Dilut to final volum of 1 lit r
- 5 g) 5 x Sample buffer Dissolve 50 mg bromophenol blue in 2.5 ml of 0.5 M Tris HCl, pH 6.8, add 4 ml 10% SDS, 2.5 ml of glycerol and store at 4°C
- h) TEMED
- 10 i) 1% Gelatin Dissolve 1 g of gelatin in 100 ml of dH₂O by warming suspension under running hot tap water
- 15 j) Gel staining solution 30% methanol, 10% acetic acid, 0.1% amido black 10 B
- k) Destaining solution 30% methanol, 10% acetic acid

Procedure

20 The procedure for preparing the gelatin zymogram used for this example utilized the following steps and reagents.

- 1) Assemble gel forming apparatus.
- 2) Prepare resolving gel solution for polymerization:

25 9% acrylamide 40 ml final volume:

in a 50 ml falcon tube
12 ml 30% acrylamide, 0.8% bisacrylamide
0.4 ml 10% SDS
30 7.5 ml 2 M Tris HCl, pH 8.8
4 ml 1% gelatin
16 ml dH₂O

- 3) After mixing, add 0.4 ml 10% ammonium persulfate and mix again.

- 4) Initiate polymerization with 40 µl of TEMED, mix and pour 32 ml of gel solution into the gel form.

- 5) Overlay solution with water saturated butanol and allow 30-45 minutes to polymeriz .

- 6) After polymerization, wash gel surface with dH₂O and allow to dry.

- 18 -

7) Prepare stacking gel solution for polymerization:

3% acrylamide 10 ml final volume:

5 to a 15 ml falcon tube add
2.5 ml 0.5 M Tris HCl, pH 6.8
0.1 ml 10% SDS
1.0 ml 30% acrylamide, 0.8% bisacrylamide
0.1 ml 10% ammonium persulfate
5.3 ml dH₂O

10 8) Initiate polymerization of stacking gel with 40 µl TEMED and mix.

9) Pour approximately 6 ml into gel form over polymerized running gel and insert well comb.

15 10) Remove air bubbles from the bottom of the comb teeth and add additional gel solution, if needed.

11) Allow stacking gel to polymerize 15-20 min.

20 12) After polymerization, remove comb from stacking gel and add 1 X electrode buffer to both electrode chambers.

13) Remove air bubbles from sample wells and bottom of gel.

25 14) Load samples and run electrophoresis at 30 mamps/gel.

15) After completion of electrophoresis, wash gel in two changes of 2.5% Triton X-100 for 60 minutes with gentle agitation at room temperature.

30 16) Discard Triton X-100 solution and place gel in 1 X collagenase buffer, incubate at 37°C for 2-4 hours or overnight at room temperature.

17) After incubation period, stain gel for 30 minutes in 0.1% amido black and then destain for approximately 90 minutes.

35 18) Zones of clearing correspond to gelatinolytic activity.

The preparation of separating gels of various concentrations are shown below.

- 19 -

		40 ml final volume			
		6%	8%	9%	10%
	30% acryl- amide, 0.8% bis	8ml	10.6ml	12ml	13.3ml
5	dH ₂ O	20ml	17.4ml	16ml	14.7ml

Soluble Collagenase Assay

The procedure for the collagenase assay of this example utilized the following preparation.

A 10 X collagenase buffer was prepared to obtain final concentrations of 0.5 M Tris, 2.0 M NaCl, 0.05 M CaCl₂, 2% Brij 35. To prepare one liter, combine:

60.55 g Tris
116.88 g NaCl
7.35 g CaCl₂
20 g Brij 35

Then dissolve in 800 ml of dH₂O and adjust the pH to 7.6 with the addition of concentrated HCl. Adjust the final volume to 1 liter and filter sterilize.

A 10 X bacterial collagenase (positive control) was prepared to obtain a final concentration of 0.5% (w/v). This involved the dissolving of 10 mg of bacterial collagenase (Sigma #C-5138) in 2 ml of 1 X collagenase buffer and storing at -20°C in 100 ml aliquots.

A bovine serum albumin (carrier protein) solution was prepared to obtain a final concentration of 0.5% (w/v) bovine serum albumin in 1 X collagenase buffer. This involved the dissolving of 100mg of bovine serum albumin in 20 ml of 1 X collagenase buffer and storing aliquoted in 1 ml fractions at -20°C.

A trichloroacetic acid-tannic acid-proline solution (TCATAP) was prepared to obtain a final concentration of 10% TCA, 0.5% tannic acid, 2 mM proline. This involved the combining of 10 ml of 100% trichloroacetic acid solution, 10 ml of 5% tannic acid solution and 1 ml of 200 mM proline solution, diluting

- 20 -

to 100 ml final volum and stor at 4°C, and replacing this solution every 4 weeks.

Procedure

- 5 1. Place enzyme sample activator (usually 1 mM pAPMA) and test solution in 1.5 ml Eppendorf tube. Combined volume must equal 60 µl. Add 1 X collagenase buffer, if necessary. If necessary, preincubate.
- 10 2. Prepare type IV collagen substrate by diluting stock ³H-type IV collagen (NEN #NET-931 lot #2511-018) 1:120 with 1 X collagenase buffer. Heat to 55°C for 10 minutes and cool on ice.
- 15 3. Add 5 µl of diluted ³H-type IV collagen solution to each assay tube. Vortex to mix and incubate at 28°C for 4 hours.
4. At the end of the incubation period, add 2 µl of carrier BSA solution and 7 µl of TCATAP solution. Vortex mix and allow to stand on ice for at least 10 minutes.
- 20 5. Pellet the precipitate by centrifuging in the microfuge for 10 minutes with the speed set at 6. Orient the tubes when placing them in the microfuge in order that the position of the pellet will be known.
- 25 6. Immediately after centrifugation, aspirate 55 µl of the supernatant and place in a scintillation vial. Add 5 ml of scintillation cocktail, shake well and count.

EXAMPLE 2

30 Cysteine Containing Peptide Inhibitors

The amino terminal sequence of A2058 melanoma cell type IV procollagenase (residues 1-17, APSIIKFPGDVAPKTD) as well as that of an internal domain (residues 472-490, DKPMGPLLIVATFNPELPEK) were synth sized f r use in preparation of monospecific antibodies. These peptides wer chosen becaus they w re obtained in th direct sequencing of th enzyme

- 21 -

(Collier, I.E., Wilhelm, S.M., Eisen, A.Z., Marmer, B.L., Grant, G.A., Seltzer, J.L., Kronberger, A., He., C., Bauer, E.A., and Goldberg, G.I., 1988, J. Biol. Chem 263, 6579-6587; Hoyhtya, M., Turpeenniemi-Hujanen, T., Stetler-Stevenson, W., Kruttsch, H., Tryggvason, K., and Liotta, L.A., 1988, FEBS Letters 233, 109-113), were confirmed in the predicted sequence from the cDNA clone (Collier, I.E., Wilhelm, S.M., Eisen, A.Z., Marmer, B.L., Grant, G.A., Seltzer, J.L., Kronberger, A., He., C., Bauer, E.A., and Goldberg, G.I., 1988, J. Biol. Chem 263, 6579-6587), and are derived from regions which do not show homology with the other metalloproteinases. The affinity purified antibodies were characterized using direct ELISA as well as competition experiments. The antibodies showed no cross reactivity with bovine serum albumin or unrelated peptides. The affinity purified antibodies are capable of immunoprecipitating the type IV procollagenase (Hoyhtya, M., Turpeenniemi-Hujanen, T., Stetler-Stevenson, W., Kruttsch, H., Tryggvason, K., and Liotta, L.A., 1988, FEBS Letters 233, 109-113). Western blots demonstrated that both antibodies recognized the type IV procollagenase as a single band in A2058 conditioned media identical to the Western blotting with the purified type IV collagenase.

The time course for pAPMA activation was followed using gelatin zymogram analysis, type IV collagenase assays and both affinity purified antibodies on immunoblots (Figure 7). Gelatin zymogram (Figure 7) analysis of A2058 melanoma cell type IV collagenase revealed a single band of gelatinolytic activity with a molecular weight of 70 kDa daltons. Incubation at 37°C in the presence of 1 mM pAPMA resulted in gradual conversion of this band of gelatinolytic activity to a lower molecular mass of 62 kDa. This conversion was completely inhibited in

- 22 -

the presence of 10 mM EDTA and did not occur in the absence of added PAPMA (not shown). Type IV collagenase assays of the purified type IV procollagenase revealed no collagenolytic activity in the absence of the organomercurial compound PAPMA during the incubation period. The enzyme could be activated by preincubation with PAPMA. The time course of activation during preincubation with PAPMA as measured by collagenase assay shows that full collagenolytic activity is obtained rapidly (Figure 8). Antibody A472-490 demonstrated a time dependent reduction in molecular weight with incubation that corresponds to that seen in the gelatin zymograms (Figure 7). This conversion appeared 50% complete by the 60 minute time point and essentially complete by the 90 minute. Antibody A1-17 corresponding to the amino terminal epitope showed a direct reduction in immunostaining during the course of PAPMA activation. These results indicate that the apparent molecular weight reduction following conversion of the latent to a stable, active collagenase with the organomercurial compound is the result of the loss of an amino terminal peptide fragment.

Gelatin-affinity purified type IV procollagenase was further purified by reverse phase HPLC both before and after PAPMA activation. The chromatograms showed essentially no change in the retention times for the latent and active forms. When the peaks were collected and analyzed by NaDodSO₄-PAGE, the procollagenase peak prior to PAPMA activation showed a single band at approximately 70 kDa under non-reducing conditions (Figure 9, insert). The collagenase peak after PAPMA activation (16 hours, 37°C) showed a single band at 62 kDa under non-reducing conditions (Figure 3A).

The material obtained from the procollagenase peak on HPLC prior to PAPMA activation was

- 23 -

subjected to direct amino acid sequencing. This material gave an amino terminal sequence that was identical to that previously determined for this enzyme (Figure 9). Direct sequencing of the pAPMA activated material, after HPLC purification, revealed a single new amino terminal sequence (Figure 9). This definitively demonstrates that pAPMA activation is accompanied by an autocatalytic removal of an amino terminal peptide fragment from the latent enzyme with a reduction in the molecular mass of about 8 kD. This cleavage occurs at only a single site as no evidence of any other amino terminal amino acids were detected upon sequencing many preparations of the purified, activated enzyme and no evidence of intermediates were detected by Western blotting or gelatin zymogram analysis.

The results of this study demonstrates that type IV collagenase is secreted in a latent proenzyme form requiring activation before obtaining collagenolytic activity. The organomercurial compound pAPMA is capable of this activation. Organomercurial activation of type IV procollagenase is accompanied by conversion of the proenzyme form to a lower molecular weight, active enzyme form by removal of an 80 amino acid residue peptide fragment from the amino terminus. Maximum collagenolytic activity is obtained rapidly following exposure to pAPMA. The attainment of this activity prior to complete conversion to the stable lower molecular weight form is consistent with a conformational rearrangement in the zymogen form that results in an active but unstable species, as has been reported for interstitial collagenase and stromelysin (Stricklin, G.P., Jeffrey, J.J., Roswit, W.T., and Eisen, A.Z., 1983, Biochemistry 22, 61-68; Murphy, G., Cockcroft, M.I., Stephens, P.E., Smith, B.J., and Docherty, A.J.P., 1987, Biochem. J. 248, 265-268). Activation occurs with highly purified type IV

- 24 -

procollagenase enzyme. Thus, the activation in the presence of the organomercurial compound is autoproteolytic since pAPMA is itself incapable of peptide bond hydrolysis. This autocatalytic mechanism of activation with organomercurial compounds has been demonstrated for other extracellular matrix degrading metalloproteinases, such as interstitial collagenase (Grant, G.A., Eisen, A.Z., Marmer, B.L., Rosweit, W.T., and Goldberg, G.I., 1987, J. Biol. Chem. **262**, 5886-5889; Whitman, S.E., Murphy, G., Angel, P., Rahmsdorf, H.-J., Smith, B.J., Lyons, A., Harris, T.J.T., Reynolds, J.J., Herrlich, P. and Docherty, A.J.P., 1986, Biochem. J. **240**, 913-916), and stromelysin (Wilhelm, S.M., Collier, I.E., Kronberger, A., Eisen, A.Z., Marmer, B.L., Grant, G.A., Bauer, E., and Goldberg, G.I., 1987, Proc. Natl. Acad. Sci. U.S.A. **84**, 6725-6729; Whitman, S.E., Murphy, G., Angel, P., Rahmsdorf, H.-J., Smith, B.J., Lyons, A., Harris, T.J.T., Reynolds, J.J., Herrlich, P. and Docherty, A.J.P., 1986, Biochem. J. **240**, 913-916; Murphy, G., Cockett, M.I., Stephens, P.E., Smith, B.J., and Docherty, A.J.P., 1987, Biochem. J. **248**, 265-268; Sanchez-Lopez, R., Nicholson, R., Gensel, M.C., Matrisian, L., and Breathnach, R., 1988, J. Biol. Chem. **263**, 11892-11899). These three metalloproteinase, type IV procollagenase, interstitial procollagenase and prostromelysin show significant homology at the amino acid level (Grant, G.A., Eisen, A.Z., Marmer, B.L., Rosweit, W.T., and Goldberg, G.I., 1987, J. Biol. Chem. **262**, 5886-5889; Collier, I.E., Wilhelm, S.M., Eisen, A.Z., Marmer, B.L., Grant, G.A., Seltzer, J.L., Kronberger, A., He, C., Bauer, E.A., and Goldberg, G.I., 1988, J. Biol. Chem. **263**, 6579-6587; Wilhelm, S.M., Collier, I.E., Kronberger, A., Eisen, A.Z., Marmer, B.L., Grant, G.A., Bauer, E., and Goldberg, G.I., 1987, Proc. Natl. Acad. Sci. U.S.A. **84**, 6725-

- 25 -

6729; Saua, J., Quinon s, S., Otani, Y., Nagase, H., Harris, E.D., Jr., and Kurkinen, M., 1988, J. Biol. Chem. 263, 6742-6745; Whitman, S.E., Murphy, G., Angel, P., Rahmsforf, H.-J., SMith, B.J., Lyons, A., Harris, T.J.T., Reynolds. J.J., Herrlich, P. and Docherty, A.J.P., 1986, Biochem. J. 240, 913-916; Sanchez-Lopez, R., Nicholson, R., Gensel, M.C., Matrisian, L., and Breatnach, R., 1988, J. Breatnach, R., 1988, J. Biol. Chem. 263, 11892-11899).

When the amino acid sequences for the amino termini of these enzymes are aligned for maximum homology (Figure 10), two correlations are observed. First, the site of autoproteolysis in type IV collagenase upon activation with PAPMA, which result in a stable, active enzyme, occurs at an identical locus to that previously reported for prostromelysin activation and the major product of interstitial procollagenase activation (Whitman, S.E., Murphy, G., Angel, P., Rahmsforf, H.-J., SMith, B.J., Lyons, A., Harris, T.J.T., Reynolds. J.J., Herrlich, P. and Docherty, A.J.P., 1986, Biochem. J. 240, 913-916; Murphy, G., Cockett, M.I., Stephens, P.E., Smith, B.J., and Docherty, A.J.P., 1987, Biochem. J. 248, 265-268). Similar sites of autoproteolysis following PAPMA treatment have been reported by others for prostromelysin (Wilhelm, S.M., Collierm, I.E., Kronberger, A., Eisen, A.Z., Marmer, B.L., Grant, G.A., Bauer, E., and Goldberg, G.I., 1987, Proc. Natl. Acad. Sci. U.S.A. 84, 6725-6729) and interstitial procollagenase (Grant, G.A., Eisen, A.Z., Marmer, B.L., Rosweit, W.T., and Goldberg, G.I., 1987, J. Biol. Chem. 262, 5886-5889). Second, the amino terminal peptide fragments which are removed during activation of all thre nzymes contain an odd number of cysteine residues. In type IV procollagenase, thr e cysteine residu s are present in the removed peptide fragment; Cys-31, Cys-36 and Cys-73. In

- 26 -

interstitial pr collagenase and prostrom lysin, there is a single cysteine residue present in the removed peptide fragment that corresponds to Cys-73 in type IV procollagenase. Thus, the conversion from an odd number of cysteine residues in the latent metalloproteinase to an even number of cysteine residues in the PAPMA activated form, appears to be a common feature in all three enzymes. The removal of an unpaired cysteine may be of functional significance. Finally, all three enzymes contain a highly conserved region immediately upstream to the activation locus consisting of the amino acid sequence PRCGVDPV. This sequence contains the unpaired cysteine residue in the propeptides of interstitial collagenase and stromelysin (Whitman, S.E., Murphy, G., Angel, P., Rahmsdorf, H.-J., Smith, B.J., Lyons, A., Harris, T.J.T., Reynolds, J.J., Herrlich, P. and Docherty, A.J.P., 1986, Biochem. J. 240, 913-916) and by homology the unpaired cysteine (out of the three present) in the type IV collagenase propeptide. A recent report has shown by site-directed mutagenesis studies of rat transin (homolog of human stromelysin), the importance of this conserved region in the auto-activation of this family of metalloproteinases. Recombinant transin forms containing mutations in this sequence showed a higher rate of spontaneous activation when compared with the native sequence (Sanchez-Lopez, R., Nicholson, R., Gensel, M.C., Matrisian, L., and Breathnach, R., 1988, J. Breathnach, R., 1988, J. Biol. Chem. 263, 11892-11899).

Therefore, the present inventors have used the gelatin zymogram technique to test a number of small peptides related to the type IV procollagenase activation locus for their ability to inhibit type IV collagenase enzyme activity. As shown in Figures 11 and 12, only those peptides incorporating the conserved region (PRCGVPDV) containing the unpaired

- 27 -

cystein were strongly inhibitory at concentration less than 0.1 mM. The cyst ine was r quir d for the activity. Table 2 shows that th minimal peptide sequence which retained inhibitor activity was actually the pentapeptide sequence RKPRC.

5

- 28 -

TABLE 2
Minimal Peptide Sequence and Sequence Requirements
for Inhibition of Type IV Collagenase.

5	<u>Peptide #</u>	<u>Sequence</u>	<u>Inhibition</u> ^(a)
	74	TMRKPRCGNPDVAN	+
	78	TMRKPRSGNPDVAN	-
	82		KPRCG-
10	85		MRKPRCG+
	86		KPRCGNP-
	88		RKPRC+
	93		RQPRC+
	94		RKARC+
15	95		QKPRC-
	101		RKPQC-
	106		KKPRC+
	107		RKPKC+
	108		RKLRC+

(a) Inhibitory activity was assessed using the modification of the gelatin zymogram technique described for Example 1. All peptides were tested at a final concentration of 1 mg/mL.

25

The data in Table 2 further demonstrate that peptide inhibitor activity requires specific charged sequence elements flanking a critical cysteinyl residue. Specifically, the two arginyl residues of the conserved sequence appeared critical for inhibitory activity. Substitution with polar uncharged residues at these positions (R to Q) resulted in loss of activity. However, the prolyl and lysyl residues between these two arginyl residues could be replaced with glycyl residues and these peptides retained inhibitory activity. This suggests that the lysyl and prolyl residues

30

35

- 29 -

which may be important for the local conformation of this region (Sanchez-Lopez et al., 1988), do not appear to directly participate in the inhibitory interaction.

5 Thus the proenzyme fragment of type IV collagenase contains a conserved sequence which is shared with other members of the matrix metalloproteinase family. This sequence is capable of inhibiting the activated type IV collagenase. This
10 inhibitor sequence requires the presence of an unpaired cysteinyl residue within a specific flanking peptide sequence that contains two positively charged residues at positions -1 and -4 with respect to the essential cysteinyl residue. Inhibition of enzyme
15 action by this inhibitor appears to be specific in that it requires two elements for inhibitor activity, an unpaired cysteinyl residue within a defined peptide sequence.

20 These data support the novel hypothesis that the RKPRC sequence which is present in the proenzyme fragment acts as an endogenous inhibitor of the enzyme through interaction of the cysteinyl residue with the metal atom coordinated at the active site. Furthermore, the sequence that surrounds the unpaired
25 cysteinyl residue appears to promote this inhibitory interaction as amino acid substitutions at positions adjacent to the cysteinyl residue can abolish the ability of the peptides to block type IV collagenolytic activity.

30 These results can be integrated in a unifying hypothesis for the structure of the latent collagenase gene family enzymes and their activation. The zymogen forms are the result of the proenzyme segment containing the conserved sequence
35 MRKPRCGN(V)PDV interacting with the enzyme active site to block catalytic activity. This interaction is stabilized through coordination of the zinc atom with

- 30 -

th unpaired sulfhydryl side chain of th cysteinyl
residu . Thus, any agent which disturbs th
cysteinyl-metal atom coordination can result in enzyme
activation and subsequent autocatalytic removal of the
proenzyme fragment. This can be accomplished by
reagents which would directly compete for the metal
atomsulfhydryl coordination, such as the organo-
mercurial reagents. It might also be accomplished by
chaotropic agents which would disturb the conformation
of the proenzyme fragment surrounding the critical
cysteinyl residue resulting in disruption of the metal
atom coordination. Nonproteolytic tissue activators
such as that described by Tyree et al., 1981 (Archiv.
Biochem. Biophys. 208, 440-443) may also activate
these enzymes by a similar mechanism of induced
conformational rearrangement, as has been proposed.

An additional preferred embodiment of the
present invention is exemplified by the propeptide
sequence 1-80 of the type IV procollagenase enzyme,
which contains the intrinsic inhibitor sequence RKPRC
incorporating an unpaired cysteine residue. In addi-
tion, the amino terminal end of this 80 amino acid
peptide contains an internal disulfide bond, and this
peptide also contains a tumor cell surface binding
site. The 80mer peptide, made either by synthetic
chemistry or by recombinant means, for example, may
therefore have advantages as an inhibitor for at least
three reasons. Firstly, its inhibitory activity on a
molar basis can be higher than a peptide because the
secondary structure of the inhibitory sequence is
accurately reproduced. Secondly the recombinant 80mer
protein has greater stability (longer half life) in
vivo due to the presence of the disulfide bond at the
amin terminus. Thirdly, sinc th 80m r contains a
c ll binding sit it can target to th surface of the
tumor cells wh re th delet rious proteolysis vent is
taking plac . Figure 13 illustrates the nucleotid

- 31 -

5 s qu nce of th portion of a human cDNA clone for
human type IV procollag nase which encod s the 80
amino acid sequence of the amino terminal peptide that
is cleaved from human type IV procollagenase upon
10 activation. A DNA segment having this nucleotide
sequence (or any other sequence specifying the same
amino acid sequence according to the universal genetic
code) can be used in various recombinant protein
production systems to produce the 80mer for use as an
15 inhibitor of matrix metalloproteinases. Alterna-
tively, the DNA segment having this nucleotide
sequence, which is derived from the human genome, can
be delivered into a mammal as part of a recombinant
expression vector to provide an internal genetic
source of production of the 80mer inhibitor of matrix
metalloproteinases.

Methods

Culture Methods

20 Human A2058 melanoma cells were grown to 80%
confluence in DMEM with 10% fetal bovine serum. The
media was then replaced with serum-free DMEM and the
culture continued for 24 hours. The serum-free condi-
tioned media was collected and concentrated by ultra-
25 filtration (Amicon YM 30 membrane) prior to storage at
-20°C.

Purification of Type IV Procollagenase

30 Type IV procollagenase was purified directly
from human A2058 melanoma cell concentrated condi-
tioned media by gelatin-Sepharose (Sigma) affinity
chromatography in 0.05 M Tris HCl, 0.005 M CaCl₂, 0.5
M NaCl, pH 7.6 buffer (TCS buffer) containing 0.02%
Brij 35 (Sigma). The enzyme was eluted using TCS
buffer containing 0.02% Brij plus 7% dimethyl
sulfoxid . The sample was then concentrated and
35 stored in the same buffer at -70°C until us . Type IV
procollag nas was further purified by r v rse-phas
HPLC prior t amino acid sequenc analysis on a Dionex

- 32 -

A1400 system equipped with a 0.46 X 10 cm RP300 column (Pierce Co.) equilibrated in 0.1% trifluoroacetic acid. The column was eluted with a linear gradient to a 60% acetonitrile.

5 Preparation of Antibodies to Synthetic Peptides

10 The peptides used in the immunization procedures were synthesized on a Biosearch 9600 peptide synthesizer. Antibodies were prepared and purified as described for Example 3. The antibody preparations were characterized by ELISA using a commercial ELISA kit (Kirkegaard and Perry Laboratories) Immulon 2 plates (Dynatech, Inc.) as shown in Figure 3.

15 Activation of Type IV Procollagenase by Organomercurials

20 Stock solutions of 0.01 M p-APMA in 0.05 N NaOH were prepared fresh daily. Proenzyme samples were incubated with a final concentration 0.5 or 1.0 mM p-APMA for varying times (0-16 hours) at 37°C. Following incubation, the samples were analyzed directly by NaDodSO₄-PAGE on 9% acrylamide gels containing 0.1% gelatin (gelatin zymogram). Alternatively, the samples were run on 9% NaDodSO₄-PAGE and electroblotted onto Immobilon P membranes (Millipore).

25 Assays for Type IV Collagenolytic Activity

30 Type IV collagenase activity was assayed in the presence of inhibitory peptides as described for Example 1. The substrate used was ³H-propionylated, human type IV collagen (New England Nuclear). The reactions were carried out at 28°C for 4 to 16 hours.

35 EXAMPLE 3

Antibodies Recognizing Metalloproteinase Peptides

 Synthetic peptides corresponding to a series of domains of Type IV procollagenase extending from the amino terminus (residues 1-17), to an internal domain near the carboxy terminus (residues 472-490) (Figure 1, Table 3) were used as antigens to generate affinity purified polyclonal antibodies which recognize

- 33 -

nized their respective domains on the native type IV procollagenase.

TABLE 3

Synthetic Peptides Used as Antigens
for Matrix Metalloproteinase (N→C)

5	1.	Ala-Pro-Ser-Pro-Ile-Ile-Lys-Phe-Pro-Gly-Asp-Ala-Pro-Lys-Thr-Asp-Lys
	2.	Glu-Leu-Ala-Val-Gln-Tyr-Leu-Asn-Thr-Phe-Tyr-Gly-Cys-Pro-Lys
10	3.	Asn-Thr-Phe-Tyr-Gly-Cys-Pro-Lys-Glu-Ser-Cys-Asn-Leu-Phe-Val-Leu-Lys
	4.	Glu-Ser-Cys-Asn-Leu-Phe-Val-Leu-Lys-Asp-Thr-Leu-Lys-Met-Gln-Lys
15	5.	Phe-Phe-Gly-Leu-Pro-Gln-Thr-Gly-Asp-Leu-Asp-Gln-Asn-Thr-Ile-Glu
	6.	Thr-Met-Arg-Lys-Pro-Arg-Cys-Gly-Asn-Pro-Asp-Val-Ala-Asn
	7.	Thr-Met-Arg-Lys-Pro-Arg-Cys-Gly-Asn-Pro-Asp-Val-Ala-Asn-Tyr-Asn-Phe-Phe-Pro-Arg-Lys-Pro-Lys
20	8.	Arg-Lys-Pro-Arg-Cys-Gly-Asn
	9.	Asn-Pro-Asp-Val-Ala-Asn-Tyr-Asn-Phe-Phe-Pro-Arg-Lys-Pro-Lys-Trp-Asp-Lys-Asn-Gin
	10.	Met-Ile-Asn-Phe-Gly-Arg-Trp-Glu-His-Gly
	11.	Lys-Tyr-Gly-Phe-Cys-Pro-Glu-Thr-Ala
25	12.	Met-Ser-Thr-Val-Gly-Gly-Asn-Ser-Glu-Gly-Ala
	13.	Met-Trp-Cys-Ala-Thr-Thr-Ala-Asn-Tyr-Asp-Asp-Arg-Lys-Trp-Gly-Phe-Cys-Pro-Asp-Gln-Gly-Tyr-Ser-Leu
	14.	Val-Ala-Ala-His-Glu-Phe-Gly-His-Ala-Met-Gly-Leu-Glu-His-Ser-Gin
30	15.	Ala-Ala-His-Glu
	16.	Asp-Lys-Pro-Met-Gly-Pro-Leu-Leu-Val-Ala-Thr-Phe-Trp-Pro-Gln-L u-Pro-Glu-Lys

- 35 -

Enzyme-linked immunosorbent assays (ELISA) were used to demonstrate that antibody binding to solid phase synthetic peptides could be competed by solution phase peptides, and that each affinity purified antibody was monospecific (Figure 3). Western immunoblotting studies of the time course of organomercurial activation process demonstrated that the antibodies recognized the solid phase enzyme in purified form, or among a complex mixture of proteins secreted by tumor cells in culture (Figure 3). Western blotting also indicated a direct loss of the amino terminal domain (residues 1-80) during the conversion to the lower molecular weight form (Figure 7). Thus, antibodies which recognize peptide domains in the first 1-80 amino terminal residues could be used to distinguish the latent from the active form of the enzyme. Antibodies recognizing peptides in Table 3 were demonstrated to be useful in solid phase or solution phase direct or competition immunoassays to detect type IV collagenase antigen in human serum and human urine. Such body fluid assays are useful for diagnosis of localized or metastatic cancer. The anti-peptide antibodies were also demonstrated to be useful in the diagnosis of human colon carcinoma by immunohistology (Table 4).

Table 4 is a summary of an immunohistology case review of human colon cancer cases demonstrating that antibodies directed against the peptides in Table 3 can be used to identify enzyme antigen associated with malignant tumor cells.

- 36 -

TABLE 4
TYPE IV COLLAGENASE IMMUNOREACTIVITY
ANTI-PEPTIDE ANTIBODY

5	TISSUE	POSITIVE/TOTAL
	<u>NORMAL GASTRIC MUCOSA</u>	0/20
	<u>GASTRIC CARCINOMA</u>	
	CONFINED TO MUCOSA	18/20*
10	INVADING THE SUBMUCOSA	20/20* *
	INVADING FULL THICKNESS	20/20* *
	<u>NORMAL COLORECTAL MUCOSA</u>	0/10
	<u>NEOPLASTIC COLONIC POLYPS</u>	1/10
	<u>COLORECTAL CARCINOMA</u>	
15	CONFINED TO MUCOSA	1/10 *
	INVADING THE SUBMUCOSA	8/10 * *
	INVADING FULL THICKNESS	18/20 * *
	<hr/>	
	* 10-30% TUMOR CELL REACTIVITY	
20	* * 50-80% TUMOR CELL REACTIVITY	

Methods

Protein Peptides

25 Synthesis of the protein peptides was carried out on a Biosearch Model 9600 peptide synthesizer using standard Merrifield solid-phase peptide synthesis protocols. The primary sequence synthesized was VAAHEFGHAMGLEHSQ which corresponds exactly to residues 371-386 in human type IV

30 collagenase as shown in Figure 1. The three His and the two Glu residues were replaced in various combinations with Ala residues and the effect of this substitution on the type IV collagenase and zymogram gelatinase activity was investigated.

35 Preparation of Antipeptide Antibodies

 Th preparation of antipeptid antibodi s utiliz d th following steps.

- 37 -

Conjugation of the peptide to BSA: To make the peptide antigenic, it must be covalently bound to BSA or another antigenic protein. To 2 mg of peptide is added one ml of PBS, and to 6 mg of BSA is added 4 ml of PBS. These solutions are combined and 5 ml of 0.25% glutaraldehyde solution is then added to this mixture. The resulting solution is stirred at room temperature for 4 hours, then dialyzed against 1 liter of PBS overnight at room temperature. The next day, the solution is concentrated to 6 ml and dispensed in 1 ml aliquots for immunization.

Immunization of rabbits: To ensure as much as possible high titers of antibody, all immunizations should be carried out using complete Freund's adjuvant. For the first two injections, 1 ml of BSA-peptide conjugate and 1 ml of CFA are emulsified and two rabbits are injected s.c. on the back with 1 ml of emulsion at approximately 30 sites. After the second injection, 0.5 ml of conjugate solution is diluted with 0.5 ml of PBS when making the emulsion. Immunizations are done every two weeks; a bleeding is done prior to immunization, and against the weeks between injections, beginning with the third or fourth injection.

Preparation of peptide antibody affinity resin: Once the anti-peptide antibody serum becomes available, the next step is to do an affinity purification step. In the first step, a peptide affinity resin is prepared. Approximately 10-12 ml of Affi-Gel 10 (BioRad Co.) is quickly washed 3X with 40 ml of cold PBS, and resuspended in cold PBS to give a total volume of about 20 ml. At the same time, 2 mg of peptide is dissolved in 1 ml of PBS, and added to the Affi-Gel suspension with immediate mixing. The resulting mixture is gently agitated overnight in the cold. The next day, sufficient 1 M Tris HCl, pH 8.0, is added to make a 0.2 M solution, and the gel

- 38 -

agitated an additional 4 hours in the cold. The gel is ready for antibody adsorption after washing 3X with 40 ml of PBS.

5 Affinity purification of antibody: The antibody-containing serum is heated at 56°C for 30 minutes, cooled and mixed with the peptide resin, which was previously allowed to settle and had excess PBS poured off. After gentle agitation overnight in the cold, the gel suspension is poured into a column
10 and washed with two gel volumes of cold 1 M acetic acid followed by 1 gel volume of PBS. The antibody eluate is taken to approximately pH 7.0 with 6 N NaOH, and the resulting solution is diafiltered (YM-30) concentrated to about 5 ml, with buffer exchanged to PBS.

15 Coupling antibody to resin: The procedure is much the same as above, except that antibody coupling is allowed to go only 1 hour before Tris is added to stop the reaction and cap up unused active sites.

20 Immunoassays: Enzyme-linked immunosorbent assays, Western blotting studies, and immunohistology was conducted using standardized, well-accepted methods.

25 Polyclonal or monoclonal antibodies to the peptides chosen and purified as described above can be labeled with suitable radioactive, enzymatic or fluorescent labels by conventional methods, which should be apparent to those skilled in the art. Immunologic assays employing peptides and antibodies
30 described herein can be applied to biologic samples of any type including body fluids, tissue extracts, or tissue sections, using conventional immunologic methods and with the aid of unlabeled, bound, or unbound antibodies or peptides. Antibodies or
35 peptides can be coupled to suitable solid phase supports such as micro-titer wells. The described anti-peptide antibodies for type IV collagenase have

- 39 -

significant advantages over antibodies made against native whole enzyme (Tryggvason, K. and Liotta, L.A. United States Patent 4,677,058). Firstly, they recognize specific domains unique to type IV collagenase which are not homologous to other prevalent metalloproteinases. This is non-obvious and overcomes a significant problem in that large proportions of the amino acid sequence of type IV collagenase is highly homologous or even identical with other metalloproteinases as shown in Figure 2. Antibodies made against the peptides in Table 3 distinguish type IV collagenase from other metalloproteinases and furthermore can distinguish activated from latent enzyme. This latter feature is very important because pathologic situations could exist in which the ratio of latent to activated enzyme is the determining factor in disease diagnosis or prognosis.

The foregoing invention has been described in some detail for purposes of clarity and understanding. It will also be obvious to one skilled in the art that various changes and combinations in form and detail can be made without departing from the scope of the invention. In particular, it is obvious to those skilled in the art that modifications in the flanking sequences of the peptide inhibitors described herein may alter stability, activity, or specificity toward individual members of the metalloproteinase family. For example, the choice of the specific sequences to the left of the activation associated cleavage site indicated in Figure 10 may cause the inhibitor to preferentially inhibit stromelysin or type I collagenase compared to type IV collagenase. This is because this region of the sequence shows some variability between different matrix metalloproteinase types. It is further obvious that certain substitutions of amino acid choices in non-critical regions of the peptides may not significantly alter the

- 40 -

inhibitory properties of the peptide. In the case of synthetic peptides to be used as immunogens or antigens for antibodies specific for a matrix metalloproteinase, one skilled in the art will understand that a unique amino acid sequence in terms of recognition by an antibody binding site consists of a sequence of from four to six amino acids which is not known to exist in another protein in the environment for which the antibody is to be used (e.g., human biological specimens. Finally, it is known in the art that recombinant protein peptides can have activity comparable to natural peptides or synthetic peptides. The embodiments of the invention can therefore obviously be produced using suitable DNA segments inserted into appropriate expression vectors as well by synthetic peptide chemistry or isolation from natural sources where available. In any case, the peptides of this invention may be purified according to a variety of standard methods well known in the art.

* * *

For purposes of completing the background description and present disclosure, each of the published articles, patents and patent applications heretofore identified in this specification are hereby incorporated by reference into the specification.

- 41 -

WHAT IS CLAIMED IS:

1. A peptide consisting essentially of an amino acid sequence having the formula $aa_1-aa_2-aa_3-aa_4-C$, wherein:

aa_1 is a basic amino acid selected from the group consisting of R and K;

aa_2 is a polar amino acid selected from the group consisting of K, Q and N;

aa_3 is a nonpolar amino acid selected from the group consisting of P, A, G, L, I and V;

aa_4 is a basic amino acid selected from the group consisting of R or K; and

C is a cysteine having a free sulfhydryl group;

said peptide further having the ability to inhibit a matrix metalloproteinase.

2. The peptide of claim 1 wherein said amino acid sequence is RKPRC.

3. The peptide of claim 1 wherein said amino acid sequence is selected from the group consisting of TMRKPRCGNPDVAN, MRKPRCG, RQPRC, RKARC, KKPRC, RKPKC, and RKLRC.

4. A purified peptide that inhibits a matrix metalloproteinase having at least a unique portion of the amino acid sequence APSPIIKFPGDVAPKTDKELAVQYLNTFYGCPKESCNLFVLKDTLKKMQ-KFFGLPQTGDLDQNTIETMRKPRCGNPDVANYNFFP.

5. The peptide according to claim 4 having a binding site specific for tumor cells.

6. The peptide of claim 4 which contains at least five amino acids.

7. A peptide consisting essentially of the amino acid sequence AAHE or the sequence VAHE, said peptide further having the ability to inhibit a matrix metalloproteinase.

- 42 -

8. A peptide of claim 7 consisting of an amino acid sequence selected from the group consisting of:

VAAHEFGHAMGLEHSQ, VAAHEFGAAMGLEHSQ,
VAAHELGHSLGLSHST, VAAHEIGHSLGLFHSA,
VVAHELTHAVTDYTAG and AAHEEICTTNEGVM.

9. A composition of matter which comprises an amount of said peptide of claim 1 or of claim 4 sufficient to inhibit a matrix metalloproteinase in a pharmaceutically acceptable carrier.

10. The composition of claim 9 which is formulated for inhalation.

11. The composition of claim 9 which is a lyophilized powder.

12. The composition of claim 9 which is an inclusion complex adapted for buccal or sublingual administration.

13. A purified peptide having the amino acid sequence of a unique portion of a matrix metalloproteinase zymogen molecule, provided that said peptide does not have the amino acid sequence of either the active metalloproteinase or the prosequence that is cleaved from the zymogen to produce said active metalloproteinase.

14. The peptide of claim 13 which contains an amino acid sequence of at least five amino acids found within the sequence numbered 1-17 of said matrix metalloproteinase zymogen molecule.

15. The peptide of claim 14 having the amino acid sequence APSPIIKFPGDVAPKTD.

16. The peptide of claim 13 which contains an amino acid sequence of at least five amino acids found within the sequence numbered 472-490 of the matrix metalloproteinase zymogen molecule.

17. The peptide of claim 16 having the amino acid sequence DKPMGPLLIVATFNPELPEK.

- 43 -

18. An antibody elicited in response to administration of a peptide of claim 13.

19. The antibody according to claim 18 having the ability to inhibit a matrix metalloproteinase.

20. An antibody elicited in response to administration of a peptide of claim 14.

21. An antibody elicited in response to administration of a peptide of claim 15.

22. An antibody elicited in response to administration of a peptide of claim 16.

23. An antibody elicited in response to administration of a peptide of claim 17.

24. A composition of matter which comprises said peptide of claim 13 in an amount sufficient to elicit an immune response together with a pharmaceutically acceptable carrier.

25. The composition of matter of claim 24 in which said peptide is covalently linked to an immunogenic carrier protein.

26. A composition of matter which comprises said antibody of claim 18 attached to a solid support.

27. A composition of matter which comprises said antibody of claim 19 in an amount sufficient to inhibit a matrix metalloproteinase in a pharmaceutically acceptable carrier.

28. A composition of matter which is at least one antibody of claim 18 attached to a solid support.

29. The composition of matter of claim 28 wherein said solid support is a microtiter plate.

30. A DNA segment wherein said segment encodes said peptide of claim 1.

31. The DNA segment according to claim 27 having the nucleotide sequence shown in Figure 13 or a unique portion thereof.

- 44 -

32. A recombinant DNA molecule comprising said DNA segment according to claim 30 and a vector.

33. A culture of cells transformed with said recombinant DNA molecule according to claim 32.

34. A method of inhibiting a matrix metalloproteinase in a mammal comprising administration of a matrix metalloproteinase-inhibiting amount of a composition of claim 9.

35. A method of treating a mammal for a degenerative process arising from activity of a matrix metalloproteinase comprising administration of a matrix metalloproteinase-inhibiting amount of a composition of claim 9.

36. The method of claim 35 wherein the matrix metalloproteinase-inhibiting composition is administered sublingually or buccally.

37. The method of claim 35 wherein the degenerative process occurs in the respiratory tract and the matrix metalloproteinase-inhibiting composition is administered by inhalation.

38. The method of claim 35 wherein the matrix metalloproteinase-inhibiting composition is administered as an intravenous infusion.

39. A method of inhibiting a matrix metalloproteinase in a mammal comprising administration of a matrix metalloproteinase-inhibiting amount of said peptide according to claim 1 or claim 4, wherein the matrix metalloproteinase-inhibiting peptide is administered by means of administering said recombinant DNA molecule according to claim 30, whereby said DNA molecule is expressed to produce said peptide.

40. A DNA segment wherein said segment encodes said peptide of claim 4.

1/13

TYPE IV PROCOLLAGENASE AMINO ACID SEQUENCE

1 A P S P I I K F P G D V A P K T D K E L A V Q Y L N T F Y G
 31 C P K E S C N L F V L K D T L K K M Q K F F G L P Q T G D L
 61 D Q N T I E T M R K P R C G N P D V A N Y N F F P R K P K W
 91 D K N Q I T Y R I G Y T P D L D P E T V D D A F A F A F Q
 121 V W S D V T P L R F S R I H D G E A D I M I N F G R W E H G
 151 D G Y P F D G K D G L L A H A F A P G T G V G G D S H F D D
 181 D E L W T L G E G Q V V R V K Y G N A D G E Y C K F F E L F
 211 N G K E Y N S C T D T G R S D G F L W C S T Y N F F E K D G
 241 K Y G F C P H E A L F T M G G N A E G Q P C K F P F E R Q G
 271 T S Y D S C T E G R T D G Y R W C G T S E D Y D R D K K Y
 301 G F C P E T A M S T V G G N S E G A P C V F F T F L G N K
 331 Y E S C T S A G R S D G L M W C A T A N Y D D D R K W G F
 361 C P D Q G Y S L F L V A A H E F G H A M G L E H S Q D P G A
 391 L M A P I Y T Y T K N F R L S Q D D I K G I Q E I Y G A S P
 421 D I D L G T G P T P T L G P V T P E I C L Q D I V F D G I A
 451 Q U R G E I F F F K D R F I W R T V T P R D K P M G P L L V
 481 A T F W P E L P E K I D A V Y E A P Q E E K A V F F A G N E
 511 Y W I Y S A S K N K T Y I F A G D K F W R Y N E V K K M D
 541 A A F N W S L I A D A W N A I P D N L D A V V D L Q G G H S
 571 P G F P K L Y L K L E N Q S L K S V K F G S I K S D W L G
 601 Y F F K G A Y Y L K L E N Q S L K S V K F G S I K S D W L G
 631 C

FIG. 1

2/13

- 1 a1-17 APSPIIKFPGDVAPKTDKEL
- 2 MBD a370-386 VAAHEFGHAMGLEHSQ
- 3 a472-490 DKPMGPLLVAATFWPELPEKID










FIG. 2

3 / 13





5/13

MBD SYNTHETIC PEPTIDE INHIBITORS

026 VAAHEFGHAMGLEHSQ	ACTIVE
038 VAAAEFGAAMGLEASQ	NOT ACTIVE
041 VAAHEFGAAMGLEHSQ	ACTIVE
039 VAAHQFGHAMGLQHSQ	PARTIAL ACTIVITY

GELATIN ZYMOGRAM 1MG/ML 1HR

NON ACTIVE PEPTIDE 038

ACTIVE PEPTIDE 026




FIG. 5

6/13





FIG. 6

7/13


FIG. 7

A


GELATIN ZYMOGRAM



Antibody A472-490



Antibody A1-17



8/13




FIG. 8

9/13

LATENT: A P S P I I K F P G D V A P

ACTIVATED: Y N F F P R K P K W D K N Q


CLEAVAGE SITE: P D V A N Y N F F P R K
80 81

FIG. 9

1 10 20 30 40 50
 APSP I I K F P G D V A P K T D K E L A V Q - Y L N T F Y G C P K E S C N L F V L - K D T - - - - - L K K - - - - M Q K
 F P A T L E T Q E Q D V D L V Q K Y L E K Y Y N L K N D G R Q V E K R - R N S G P V V E K L K - - - - Q M Q E
 Y P L D G A A R G E D T S M N L V Q K Y L E N Y Y D L E K D V K Q - F V R R K D S G P V V - - - - K K I R E M Q K

10/13

60 70 80 90 100 110
 F F G L P Q T G D L D Q N T I E T M R K P R C G N P D V A N Y N F F P R K P K W D K N Q I T Y R I I G Y T P D L D P E T
 F F G L K V T G K P D A E T L K V M K Q P R C G V P D V A Q F V L T E G N P R W E Q T H L R Y R I E N Y T P D L P R A D
 F L G L E V T G K L D S D T L E V M R K P R C G V P D V G H F R T F P G I P K W R K T H L T Y R I V N Y T P D L P K D A

A B

FIG. 10

11/13




FIG.11

1 10 20 30 40 50
 APSPIIKFPGDVAPKTDKELAVQ--YLNTFYGCPKESCNLFV L-KDT-----LKK---MQK

APSPIIKFPGDVAPKTDK NOT ACTIVE

ELAVQYLNTFYGCPK NOT ACTIVE

60 70 80 90 100 110
 FFGLPQTGDL DQNTIE TMRKPRCGNPDVANYNFFPRKPKWDKNQITYRIIGYTPDLDPET

12/13

NPDVANYNFFPRKPKWDKNQ NOT ACTIVE

TMRKPRCGNPDVANYNFFPRKPK ACTIVE

TMRKPRCGNPDVAN ACTIVE

FIG. 12

SEQUENCE RANGE: 1 TO 240

GCG CCG TCG CCC ATC ATC AAG TTC CCC GGC GAT GTC GCC CCC AAA ACG GAC
 A P S P I I K F P G D V A P K T D>

100
 *
 AAA GAG TTG GCA GTG CAA TAC CTG AAC ACC TTC TAT GGC TGC CCC AAG GAG
 K E L A V Q Y L N T F Y G C P K E>
 13/13
 AGC TGC AAC CTG TTT GTG CTG AAG GAC ACA CTA AAG AAG ATG CAG AAG TTC
 S C N L L F V L K D T L K K M Q K F>


200
 *
 TTT GGA CTG CCC CAG ACA GGT GAT CTT GAC CAG AAT ACC ATC GAG ACC ATG
 F G L P Q T G D L L D Q N T I E I M>

CGG AAG CCA CGC TGC GGC AAC CCA GAT GTG GCC AAC
 R K P R C G N P D V A N>

FIG. 13

INTERNATIONAL SEARCH REPORT

International Application No. **PCI/US90/01060**

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC(5): G01N 33/543, 33/00; C12N 5/00, 1/20 U.S. CL.: 436/518, 94; 435/240.27, 253		
II. FIELDS SEARCHED		
Minimum Documentation Searched ?		
Classification System	Classification Symbols	
U.S.	436/518, 94; 530/387, 300, 330, 329, 327; 435/240.27, 253	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched *		
Chemical Abstracts Services Online, Automated Patent System		
III. DOCUMENTS CONSIDERED TO BE RELEVANT *		
Category *	Citation of Document, ** with indication, where appropriate, of the relevant passages **	Relevant to Claim No. **
Y	Journal of Biological Chemistry, Volume 262, issued 1987, J. H. MCKERROW, "Human fibroblast collagenase contains amino acid sequence homologous to the zinc-binding site of Serraha protease", page 5943.	1-40
Y	Journal of Biological Chemistry, Volume 258, issued 1983, SALO, T. ET AL., "purification and characterization of a murine basement membrane collagen-degrading enzyme secreted by metastatic tumor cells", pp 3058-3063, see the results.	1-40
Y	Biochemistry, Volume 20, issued 1981, LIOTTA, L.A. ET AL., "Partial purification and characterization of a neutral protease which cleaves type IV collagen", pp. 100-104, see the results.	1-40
<div style="display: flex; justify-content: space-between;"> <div style="width: 48%;"> <p>* Special categories of cited documents: **</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 48%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
09 APRIL 1990		18 MAY 1990
International Searching Authority		Signature of Authorized Officer
ISA/US		 KAREN I. KRUPEN

III. D CUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	Biochimica et Biophysica Acta, Volume 831, issued 1985, MRUPHY, G. ET AL., "Purification and characterization of a bone metalloproteinase that degrades gelatin and types IV AND V collagen", pp 49-58, see the results.	1-40
Y	Journal of Biological Chemistry, Volume 264, issued 1989, STETLER-STEVENSON, W.G. ET AL., "The Activation of human type IV collagenes proenzyme", pp 1353-1356, see the introduction and results.	1-40
Y	FEBS Letters, Volume 233, issued 1988, HOYHIYA, M. ET AL., "Monoclonal antibodies to type IV collagenase recognize a protein with limited sequence homology to interstitial collagenase and Stromelysin", pp 109-113, see the materials and methods, p. 110 and results, pp 111-112.	18-29
Y	US, A, 4,486,530 (DAVID) 04 December 1984, see column 5, line 50 through column 8, line 3.	26,28,29
Y	Methods in Enzymology, Volume 70, issued 1980, HURN, B.A.L., ET AL, "Production of reagent antibodies", pp 104-135, see pp 105-110.	18-25